



UNIVERSITI PUTRA MALAYSIA

**PLANTLET REGENERATION AND CALLUS INDUCTION
IN HYBRID CABBAGE
(BRASSICA OLERACEA L VAR. CAPITATA CV. TROPICANA)**

TENGKU LAILA KAMALIAH

FP 2002 17

**PLANTLET REGENERATION AND CALLUS INDUCTION
IN HYBRID CABBAGE
(*BRASSICA OLERACEA* L. VAR. *CAPITATA* CV. TROPICANA)**

TENGGU LAILA KAMALIAH

**MASTER OF AGRICULTURAL SCIENCE
UNIVERSITI PUTRA MALAYSIA**

2002



**PLANTLET REGENERATION AND CALLUS INDUCTION
IN HYBRID CABBAGE
(*BRASSICA OLERACEA* L. VAR. *CAPITATA* CV. TROPICANA)**

By

TENGKU LAILA KAMALIAH

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Requirement for the Degree of
Master of Agricultural Science**

October 2002



In the name of Allah, The Most Beneficent, The Most Merciful

*Say: "Truly, my prayer and my service of sacrifice and my life and my death,
are (all) for Allah, the Cherisher of the Worlds:*

(Q.S. 6:162)

to :

My beloved husband,

Idam Kalim

My parents (K. Tengku Mustafa Kamal and Hj. Tengku Cici Nazariah)

My parents-in-law (Abdul Kalim and Fatimah Kals)

My brother (Tengku Julian Mustafizar)

mg/L BA + 30 g/L sucrose can be used for subculturing multiple shoots derived from shoot tip explants. Subculturing should be limited to three subcultures.

There were two treatment options for induction of multiple shoots from nodal explants: 5 mg/L AgNO₃ or 0.1 mg/L NAA (naphthalene acetic acid), combined with 3 mg/L BA and 45 g/L sucrose. For subculturing of multiple shoots derived from nodal explants, medium containing 2 mg/L BA + 30 g/L sucrose can be used. Subculturing should be limited to three subcultures.

The specific objectives of the study on shoot elongation, rooting and acclimatisation were: to find out suitable medium for shoot elongation, to establish suitable media type and concentration of auxin for rooting, to determine an appropriate medium for acclimatisation of micropropagated plantlets and to determine stomata characteristic and number of stomata per mm² of leaves of plantlets in test tube, under acclimatised and *ex vitro* condition. All experiments were conducted and arranged in a Completely Randomised Design (CRD), except for experiment on “Effect of Growth Media on *Ex Vitro* Performance”, which was conducted in a Randomised Complete Block Design (RCBD).

The results showed that full strength MS medium without growth regulators was suitable for shoot elongation and rooting. Soil and oil palm frond compost mixture (1:1, v/v) was a suitable medium for acclimatisation. Plantlets were well rooted and ready to be transferred to field condition after

Experiments were also conducted with the specific objectives to induce callus from cotyledon explants and to study the effect of 2,4-dichlorophenoxy acetic acid (2,4-D), kinetin and sucrose on callus formation. The best treatment for callus formation was 1 mg/L 2,4-D with 30 g/L sucrose. Higher level of 2,4-D decreased percentage of callus formation. The presence of kinetin at all concentrations tested and the addition of sucrose above 30 g/L decreased the production of callus.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains Pertanian

**REGENERASI PLANTLET DAN INDUKSI KALUS
KOBIS HIBRID
(*BRASSICA OLERACEA* L. VAR. *CAPITATA* CV. TROPICANA)**

Oleh

TENGGU LAILA KAMALIAH

Oktober 2002

Pengerusi : Azmi Abdul Rashid, M.Phil.

Fakulti : Pertanian

Penyelidikan ini dilaksanakan dengan objektif utama untuk mewujudkan sebuah protokol untuk meregenerasi anak pokok *Brassica oleracea* L. subsp. *capitata* cv. Tropicana melalui penggandaan tunas dan untuk menginduksi kalus dengan menggunakan eksplan kotiledon. Kajian ini terdiri dari beberapa bahagian iaitu penggandaan tunas; pemanjangan pucuk, pengakaran dan aklimatisasi; dan induksi kalus.

Objektif khas daripada kajian penggandaan tunas adalah untuk menentukan pengawalatur tumbesaran dan kepekatan sukrosa dan argentum nitrate (AgNO_3) yang sesuai untuk menginduksikan tunas berganda dan untuk mengkaji kesan daripada 6-benzylaminopurina (BA) terhadap penggandaan tunas. Kesemua eksperimen dilaksanakan dan disusun mengikut Rekabentuk Rawak Lengkap.

Hasil kajian menunjukkan bahawa BA lebih baik dari kinetin untuk menggalakkan pembentukan tunas berganda daripada eksplan mercu pucuk. Argentum nitrat (AgNO_3) pada kepekatan 0.5 mg/L dikombinasikan bersama 3 mg/L BA dan 45 g/L

sukrosa meningkatkan penggandaan tunas. Medium yang mengandungi 0.5 mg/L BA + 30 g/L sukrosa boleh digunakan untuk pengsubkulturan. Pengsubkulturan sebaiknya dihadkan sehingga tiga kali sahaja.

Ada dua pilihan rawatan untuk menggalakkan penggandaan tunas daripada eksplan buku iaitu: 5 mg/L AgNO₃ atau 0.1 mg/L NAA (asid naptalena asetik), dikombinasikan bersama 3 mg/L BA dan 45 g/L sukrosa. Bagi pengsubkulturan tunas daripada eksplan buku, medium yang mengandungi 2 mg/L BA + 30 g/L sukrosa boleh digunakan. Subkultur sebaiknya dihadkan sehingga tiga kali sahaja.

Objektif khas daripada kajian pemanjangan pucuk, pengakaran dan aklimatisasi adalah untuk mengetahui medium yang sesuai untuk menggalakkan pemanjangan pucuk, untuk mewujudkan jenis medium dan kepekatan auksin yang sesuai untuk pengakaran, untuk menentukan medium yang sesuai untuk mengaklimatisasi anak pokok yang dibiak secara pembiakan mikro dan untuk menentukan ciri-ciri stomata dan bilangan stomata per mm² bagi daun anak pokok tabung uji, yang telah diaklimatisasi dan ditanam dipersekitaran *ex vitro*. Kesemua eksperimen telah dilaksanakan dan disusun mengikut Rekabentuk Rawak Lengkap, kecuali eksperimen “Kesan Medium Pertumbuhan terhadap Prestasi Anak Pokok di Persekitaran *Ex Vitro*”, yang disusun mengikut Rekabentuk Rawak Berblok Penuh.

Hasil kajian menunjukkan bahawa medium MS berkepekatan penuh tanpa pengawalatur tumbesaran sesuai digunakan untuk menggalakkan pemanjangan pucuk dan pengakaran. Campuran tanah dan kompos pelepah kelapa sawit (1:1, v/v) adalah medium yang sesuai untuk aklimatisasi. Anak pokok berakar dengan baik dan

tersedia untuk dipindahkan ke kondisi ladang sesudah 4 minggu dalam kondisi aklimatisasi.

Kajian telah juga dilaksanakan dengan objektif khas untuk menginduksi kalus dari eksplan kotiledon dan untuk mengkaji kesan 2,4-asid diklorofenoksi asetik (2,4-D), kinetin dan sukrosa terhadap pembentukan kalus. Rawatan yang terbaik untuk pembentukan kalus adalah 1 mg/L 2,4-D bersama 30 g/L sukrosa. Peningkatan kepekatan 2,4-D menurunkan peratus pembentukan kalus. Kehadiran kinetin pada seluruh kepekatan yang diuji dan penambahan sukrosa melebihi 30 g/L menurunkan kadar pembentukan kalus.

ACKNOWLEDGEMENTS

All Praise and Thanks are due to Allah who gave me strength and easiness to complete my study. I would like to express my gratitude to the chairman of my supervisory committee, Azmi Abdul Rashid, M.Phil., for his help and guidance during my study. I am also indebted and very grateful to my committee member, Dr. Maheran Abdul Aziz, for her kind advice and constant support during my study. Thanks are also due to Dr. Siti Aishah Hasan for her help and suggestions and Dr. Thohirah Abdullah as chairperson viva for her suggestions.

I wish to acknowledge the invaluable assistance of Mr. Agus Sutanto and Mr. Md. Humayun Khabir in laboratory works, Mr. Budi Winarto in discussions, Miss Maiyastri in statistical analysis, Mr. Erwan in providing oil palm frond compost, Dr. Said Sa'ad and Mrs. Halimah Alma. I appreciate all members of Makmal *In Vitro*, Aini, Beverlien and Elixon for their invaluable help. A special thank you for my lecturer at Bogor Agricultural University, Dr. Sriani Sujiprihati, Prof. Dr. Sri Setyati Harjadi and Mrs. Ketty Suketi for moral support. My friends at Jakarta (Mila cs), my friends at Indonesian Student Association and a number of people have been continuously very supportive and pray for me, thanks a lot for your support and prayer.

Finally, I would like to thank my husband, my parents, my parents-in-law and my brother. Thanks are also due to my sister-in-law and others member of my family for their endless prayer, love, patience and understanding have been a great source of encouragement to me.


I certify that an Examination Committee met on 31st October 2002 to conduct the final examination of Tengku Laila Kamaliah on her Master of Agricultural Science thesis entitled “PLANTLET REGENERATION AND CALLUS INDUCTION IN HYBRID CABBAGE (*BRASSICA OLERACEA* L. VAR. *CAPITATA* CV. TROPICANA)” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) regulation 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Thohirah Abdullah, Ph.D.
Department of Crop Science,
Faculty of Agriculture,
Universiti Putra Malaysia,
(Chairperson)

Azm i Abdul Rashid, M.Phil.
Department of Crop Science,
Faculty of Agriculture,
Universiti Putra Malaysia.
(Member)

Maheran Abdul Aziz, Ph.D.
Department of Crop Science,
Faculty of Agriculture,
Universiti Putra Malaysia.
(Member)

Siti Aishah Hasan, Ph.D.
Department of Crop Science,
Faculty of Agriculture,
Universiti Putra Malaysia.
(Member)



SHAMSHER MOHAMAD RAMADILI, Ph.D.,
Professor / Deputy Dean,
School of Graduate Studies,
Universiti Putra Malaysia

Date: 13 NOV 2002

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfillment of the requirement for the degree of Master of Agricultural Science. The member of the Supervisory Committee are as follows :

Azmi Abdul Rashid, M.Phil.

Department of Crop Science,
Faculty of Agriculture,
Universiti Putra Malaysia.
(Chairman)

Maheran Abdul Aziz, Ph.D.

Department of Crop Science,
Faculty of Agriculture,
Universiti Putra Malaysia.
(Member)

Siti Aishah Hasan, Ph.D.

Department of Crop Science,
Faculty of Agriculture,
Universiti Putra Malaysia.
(Member)



AINI IDERIS, Ph.D.,
Professor / Dean,
School of Graduate Studies,
Universiti Putra Malaysia

Date : 08 JAN 2013

DECLARATION

I hereby declare that this thesis is based on my original work except quotations and citations which have been duly acknowledge. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



TENGKU LAILA KAMALIAH

Date : 12 / 11 / 2002

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVAL	x
DECLARATION	xii
LIST OF TABLES	xvi
LIST OF FIGURES	xvii
LIST OF PLATES	xix
LIST OF ABBREVIATIONS/NOTATIONS	xx
 CHAPTER	
I GENERAL INTRODUCTION	1
Background	1
Objectives	3
 II LITERATURE REVIEW	5
Botany of Cabbage	5
<i>In Vitro</i> Plantlet Regeneration	6
Axillary Shoot Proliferation	6
Adventitious Shoot Formation	8
Somatic Embryogenesis	9
Factors Affecting Axillary Shoot Proliferation and Adventitious Shoot Formation	10
Types of Explant	10
Choice of Nutrient Medium	12
Plant Growth Regulators	13
Cytokinin	13
Auxin	15
Sucrose	16
Ethylene and Silver Nitrate	18
Physical Environment of Culture	19
Factors Affecting Somatic Embryogenesis	21
Types of Explant	21
Choice of Nutrient Medium	21
Plant Growth Regulators	22
Sucrose	23
Physical Environment of Culture	23
Rooting	24
Acclimatisation	26
 III PLANTLET REGENERATION VIA INDUCTION OF MULTIPLE SHOOT	30
Preface	30



Materials and Methods	31
Explants Preparation and Culture	31
Experiments	32
Histology	35
Experimental Design and Statistical Analysis	35
Results	38
Induction of Multiple Shoots	38
Effect of BA and NAA on Multiple Shoot Formation from Cotyledon Explants	43
Effect of Different Concentrations of BA on Multiple Shoot Formation from Shoot Tip Explants	43
Effect of Different Concentrations of Kinetin on Multiple Shoot Formation from Shoot Tip Explants	46
Comparison between BA and Kinetin on Multiple Shoot Formation from Shoot Tip Explants	48
Effect of BA and Sucrose on Multiple Shoot Formation from Shoot Tip Explants	50
Effect of AgNO ₃ and NAA on Multiple Shoot Formation from Shoot Tip Explants	54
Effect of AgNO ₃ and NAA on Multiple Shoot Formation from Nodal Explants	57
Effect of Subculturing on Multiple Shoot Formation	59
Discussion	60
Multiple Shoot Formation	60
Effect of Subculturing on Multiple Shoot Formation	65

IV	SHOOT ELONGATION, ROOTING AND ACCLIMATISATION OF PLANTLET TO EX VITRO ENVIRONMENT	66
	Preface	66
	Materials and Methods	68
	Explants Preparation and Culture	68
	Experiments	69
	Experimental Design and Statistical Analysis	72
	Results	73
	Shoot Elongation	73
	Effect of MS Salt Strength and IBA Concentrations on Rooting	75
	Comparison between IBA and NAA on Rooting	78
	Effect of Growth Media on <i>Ex Vitro</i> Performance	79
	Comparison of Stomata Number and Characteristic between Plantlets from Unacclimatised (in Test Tube), under Acclimatised and Field Condition	85
	Discussion	87
	Shoot Elongation	87
	Rooting	87
	Acclimatisation	88
	Comparison of Stomata Number and Characteristic between Plantlets from Unacclimatised (in Test Tube), under Acclimatised and Field Condition	89

V	CALLUS INDUCTION	91
	Preface	91
	Materials and Methods	92
	Explants Preparation and Culture	93
	Experiments	94
	Results	95
	Effect of Different Concentrations of 2,4-D on Callus Induction	95
	Effect of Different Concentrations of Kinetin and Sucrose on Callus Induction	98
	Discussion	100
VI	GENERAL DISCUSSION AND CONCLUSION	102
	REFERENCES	107
	APPENDICES	116
	BIODATA OF THE AUTHOR	131

LIST OF TABLES

Table	Page
1 The combination of BA (B) and NAA (N) concentrations for induction of multiple shoots from cotyledon explants	36
2 The combination of BA (B) and sucrose (S) concentrations for induction of multiple shoots from shoot tip explants	36
3 The combination of AgNO ₃ (A) and NAA (N) concentrations for induction of multiple shoots from shoot tip explants	37
4 The combination of AgNO ₃ (A) and NAA (N) concentrations for induction of multiple shoots from nodal explants	37
5 Effect of BA and sucrose on multiple shoot formation	50
6 Mean number of transferable shoots per explant from nodal explants on different concentrations of AgNO ₃ and NAA after 6 weeks of culture ...	57
7 The combination of MS (M) and BA (B) concentrations for shoot elongation	73
8 The combination of MS (M) salt strength and IBA (I) concentrations for rooting formation	73
9 Shoot elongation in different MS salt strength and BA concentrations ...	74
10 Mean number of roots formed per explant on different MS salt strength and IBA concentrations	75
11 Percentage of root formation on different MS salt strength and IBA concentrations	77
12 Comparison between IBA and NAA on rooting	79
13 The combination of kinetin (K) and sucrose (S) concentrations on callus induction from cotyledon explants	95
14 Effect of different concentrations of 2,4-D on callus induction	96
15 Effect of different concentrations kinetin and sucrose on callus induction	99
16 Summary of experiments on multiple shoot formation	104
17 Summary of experiments on subculturing	105



LIST OF FIGURES

Figure	Page
1 Effects of different concentrations of BA on mean number of transferable shoots per explant and percentage of shoot formation from shoot tip explants after 7 weeks of culture	45
2 Effects of different concentrations of kinetin on mean number of transferable shoots per explant from shoot tip explants after 7 weeks of culture	46
3 Effects formation from shoot tip explants after 7 weeks of culture	47
4 Comparison between BA and kinetin on multiple shoot formation from shoot tip explants on mean number of transferable shoots per explant was taken until 7 weeks of culture	48
5 Comparison between BA and kinetin on multiple shoot formation from shoot tip explants on percentage of shoot formation after 7 weeks of culture	49
6 Effect of BA and sucrose on mean number of transferable shoot per explant and percentage of shoot formation from shoot tip explants after 7 weeks of culture	51
7 Effect of BA and sucrose on total fresh weight and total dry matter after 7 weeks of culture	52
8 Effect of AgNO ₃ and NAA on mean number of transferable shoots per explant from shoot tip explants after 7 weeks of culture	54
9 Effect of AgNO ₃ and NAA on percentage of shoot formation (%) from shoot tip explants after 7 weeks of culture	55
10 Effect of AgNO ₃ and NAA on percentage of rhizogenesis (%) from shoot tip explants after 7 weeks of culture	56
11 Effect of AgNO ₃ and NAA on percentage of rhizogenesis (%) from nodal explants after 7 weeks of culture ...	58
12 Effect of subculturing on multiple shoot formation from shoot tip explants with 3 weeks interval of subculture	59
13 Effect of subculturing on multiple shoot formation from nodal explants with 3 weeks interval of subculture	60

14	Total chlorophyll content (mg/mg) from 0 until 3 weeks after transplanting	80
15	Effect of growth media on <i>ex vitro</i> performance on number of leaves developed per explant and plant height (cm)	81
16	Effect of growth media on <i>ex vitro</i> performance on total fresh weight and total dry matter (g)	81
17	Percentage of plants survival (%)	82
18	Comparison of stomata number between plantlets from unacclimatised (in test tube), under acclimatised and field condition	85

LIST OF PLATES

Plate		Page
1	A flowchart on possible pathway of clonal propagation of white cabbage from different explants	4
2	Multiple shoot formation from shoot tip explant	40
3	Multiple shoot formation from nodal explant	41
4	Histological observation on adventitious shoot formation from callus at the base of shoot tip	42
5	Rhizogenesis occurring from cotyledon explants	43
6	Multiple shoot formation on different concentrations of BA	44
7	Multiple shoot formation on different concentrations of kinetin	47
8	Effect of BA and sucrose on multiple shoot formation	53
9	Rhizogenesis occurring on nodal explants	58
10	Rooting induction	76
11	Plant survival and plant death in the first week after transplanting	83
12	Plantlets survival in tray containing various media after three weeks of transplanting under acclimatised conditions	83
13	Plantlets survival after three weeks of transplanting in tray	84
14	Plant survival at outdoor environment (after 3 months of acclimatisation)	84
15	Stomata from leaves of <i>in vitro</i> plantlets (in test tube) are wide open ...	86
16	Stomata from leaves of plantlets under acclimatised condition are only slightly open	86
17	Callus induction	97
18	Rhizogenesis appeared at the second subculture	100



LIST OF ABBREVIATIONS/NOTATIONS

A_{647}	absorbance at 647 nanometers
A_{664}	absorbance at 664 nanometers
ACCA	1-amino-cyclopropane-1-carboxylic-acid
AgNO_3	silver nitrate
$\text{Ag}_2\text{S}_2\text{O}_3$	silver thiosulfate
2,4-D	2,4-dichlorophenoxyacetic acid
2-iP	γ,γ -(dimethylallyl) aminopurine
α	level of significance
ANOVA	analysis of variance
Arc sin	$\sin^{-1}(X)$
AVG	aminoethoxyvinylglycine
BA	6-benzylaminopurine
cm	centimeter
CRD	completely randomised design
cv.	cultivar
DNMRT	duncan new multiple range test
e.g.	exempli gratia (for example)
<i>et al.</i>	et alia
etc.	et cetera
FAA	formaldehyde-glacial acetic acid alcohol
F_1	first filial generation
Fe-EDTA	iron ethylene diamine tetra acetic acid
g	gram

g/L	gram per litre
IAA	indole acetic acid
IBA	indole butyric acid
mg	milligram
mg/L	milligram per litre
mg/mg	milligram per milligram
mL	millilitre
mm	millimetre
$\mu\text{mol m}^{-2} \text{s}^{-1}$	micromole per metre square per second
MS	Murashige and Skoog (1962) basal medium
N	nitrogen
NAA	naphthalene acetic acid
NCSS	Number Cruncher Statistical System
%	percent
pH	$\log(\text{H}^+)$
RCBD	randomised complete block design
SAM	S-adenosylmethionin
sp.	species
$\sqrt{\quad}$	square root
var	variety
v/v	volume to volume

CHAPTER 1

GENERAL INTRODUCTION

Background

Cabbage is among the important vegetable crops in the world. It is cultivated in many countries in the northern and southern hemisphere. Asia clearly leads producing 58% of the world cabbage production (Rubatzky and Yamaguchi, 1997). In tropical Asia the consumption of early maturing cabbage, with firm and round heads, is rapidly increasing.

In Malaysia, the annual consumption of cabbage per capita is the second highest in 1991 after Chinese mustard (Kamil *et al.*, 2001). Cabbage is regarded as a good source of calcium (22.8 mg/100 g fresh weight) and dietary fibre (2.5 g/100 g fresh weight) (Lucarini *et al.*, 1999). In addition, it possesses anticarcinogenic properties. Two bioactive sulfur-containing phytochemicals, glucosinolates and S-methyl cysteine sulf oxide, which is formed in cabbage appear to have potential value as cancer chemopreventive agents (Verhoeven *et al.*, 1997).

In Malaysia, seeds of *Brassica*, including cabbage, are the second most commonly imported vegetable seeds. The total amount of *Brassica* seeds imported was about 391,246 kg and exceeding RM 300,000 in 1998 (Puteh and Abidin, 1999). Most of the *Brassica* seeds imported were F1 hybrids.

Malaysia imports not only seeds of cabbage but also fresh cabbage from Indonesia and Australia. Quantity import of fresh cabbage increases from year to year. Malaysia imported 16,770.76 ton cabbage in 1980, 28,835.12 ton in 1990 and 30,596.06 ton in 1995 (Lin, 1997).

The production of hybrid seeds of cabbage is almost impossible in Malaysia because of climatic hindrance. Vernalisation is needed for the induction of flowering. Moreover, the parent material for hybridisation and production of hybrid seed in Malaysia is not available. Therefore an alternative method of producing and maintaining plants having F₁ hybrid characteristic is through *in vitro* culture. On the other hand, high regeneration capability through tissue culture system is crucial to the success of *Brassicas* genetic engineering through genetic transformation (Takasaki *et al.*, 1997).

In vitro plantlet regeneration can be achieved through axillary proliferation, adventitious shoot formation and somatic embryogenesis. Axillary proliferation can be obtained by using shoot tip culture or axillary bud culture. Adventitious shoot formation and somatic embryogenesis can be achieved directly from explants or indirectly via callus formation. Indirect somatic embryogenesis is the most promising regeneration system for future genetic improvement of cabbage. Genetic improvement of cabbage through conventional breeding is hindered by sexual incompatibility. Therefore, the development of protocol to induce embryogenic callus and somatic embryogenesis are needed to obtain pest and disease resistant cabbage in the future.

Plate I shows the possible steps of clonal propagation of white cabbage, beginning from germination of seed, multiple shoot induction, rooting, acclimatisation and field planting. In future, this protocol can also be used for cabbage genetic improvement.

Objectives

The protocol for *in vitro* plantlet regeneration of white cabbage cv. Tropicana has not been developed. In addition, regeneration of whole plants from cultured tissues or cells is a prerequisite for successful applications of *in vitro* techniques for gene transfer and mass propagation. Therefore, the objectives of this research were: (a) to establish a protocol for plantlet regeneration of *Brassica oleracea* L. var. *capitata* cv. Tropicana through multiple shoot formation and (b) to induce callus using cotyledonary explants, which if embryogenic, can be used for inducing somatic embryos.